

# The role of host genetic variability in the development and establishment of human gut microbiome diversity

#### Larissa Matos

Instituto de Biociências, Centro de Pesquisa sobre o Genoma Humano e Células-Tronco, USP

## ABSTRACT

**Background:** Studies have shown that human microbiome plays important roles in physiology, from food digestion to mental diseases. Since the gut microbiome composes the highest number of microbial cells outnumbering even our own cell counts, it is expected that the gut microbiome would allect a great deal of human biological functions. This makes the gut microbiome key to maintain homeostasis in the various biological levels where there are constant and active interactions between microbes, tissue, cells and molecules. The structure of the gut microbiome during the development and establishment of the gut microbiota at early stages of human life is crucial to infer biological and pathological microbiome composition. The purpose of this study is to apply tools of bioinformatics to process and analyse data samples of feces of triplet babies in order to verify host genetic associations with gut microbiome during the lrst 3 years of the babies lives.

**Methods:** Infant feces are collected from the list week of life until 3 years of age. DNA extraction from the samples is performed followed by PCR (Polymerase Chain Reaction) targeting the specilic bacterial 16S ribosomal RNA gene, region V3 - V4. Next-Generation sequencing is applied. Amplicons are then computationally processed by elicient algorithms to yield high quality reads. Later, these reads are assigned to groups of taxa based on dillerences on single nucleotide. Species specilic classillers and database are used for taxonomic assignment. Alpha and beta diversity are analysed with both qiime2 plugins and R packages.

**Results:** Most of our samples presented enough reads to identify all taxa. Phylogenetic diversity increased in samples on later time points presenting time as the dominant factor to determine alpha and beta diversity. Actinobacteria, Verrucomicrobia, Proteobacteria, Firmicutes and Bacteroidetes were the dominant phyla in all samples. At the species level, monozygotic twins presented more similar microbiome between them than between their dizygotic sibling.

**Conclusions:** Our results showed that genetic factors could be detected at the species level and that time is crucial for diversity in the <code>lrst</code> months of life. The identilcation of the gut microbiome structure in both monozygotic and dizygotic twins sheds some light on how the development and establishment of microbiota take place on the human gut. For our next analysis we will apply appropriate statistics for twin studies and check specilc species and its association with host genetics.

## Biography

She is is a Molecular Biologist and Academic Scientific Consultancy. She is from Brazil

#### Publication of speakers:

 Romagnolli, Lucas & Coria, Vivian & Júnior, Sérgio & Sartori, João & Romanelli, Alethea & Oliveira, Júlio & Carvalho, Lívia & Martin, Bruno & Fornasari, Marina & Avila, Lazslo & Filho, Gerardo. (2016). Alexithymia and fibromyalgia: a systematic review. Revista de Medicina. 95. 12. 10.11606/ issn.1679-9836.v95i1p12-17.

 Coria, Vivian & Grigolo, Ingrid & Mutter, Verena & Defende, Gabriela & Moras, Luis & Morales, Stephanie & Carrai, Larissa & Dias, Ana & Costa, Nauyla & Budin, Alexandre & Rodrigues, Camila & Marchiori, Renata & Sasso, Danilo & Calocini, Andre & Lima, Juliana & Reis, Andréia & Kuyumjian, Fátima & Cação, João. (2017). Caracterização dos idosos internados por doença respiratória aguda em um hospital escola terciário. Revista de Medicina. 96. 94. 10.11606/issn.1679-9836.v96i2p94-102.

**Citation:** Larissa Matos ; The role of host genetic variability in the development and establishment of human gut microbiome diversity; Clinical Microbiology 2021; London, UK

**Copyright:** © 2021 Matos L. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.